



Visit cvs.co/ai-list or scan the QR code to view electronically.
For easy access, you can bookmark this page.

Autoimmune (AI) Drug List

Select FDA-approved AI specialty medications and infusion therapy

Drug name	Manufacturer	FDA-Approved Indication(s)	Route of Administration	Supply Method	Storage	Select Common Adverse Reactions
Actemra (tocilizumab)	Genentech	• CRS (2+) • GCA • PJIA (2+) • RA • SJIA (2+) • SSc-ILD • Hospitalized adults who have COVID-19	SC injection: • Self-administered IV infusion: • Prepared and infused by a health care professional	SC: • 162 mg/0.9 mL PFS • 162 mg/0.9 mL PEN IV: • 80 mg/4 mL • 200 mg/10 mL • 400 mg/20 mL	• Refrigerate at 36°F to 46°F (2°C to 8°C) • Do not freeze • Store in original packages • Protect from light • Keep syringes dry	Upper respiratory tract infections, nasopharyngitis, headache, hypertension, increased alanine aminotransferase (ALT) and injection site reactions
Adalimumab-aacf (unbranded version of Idacio)	Fresenius Kabi	• AS • CD (6+) • HS • PJIA (2+) • PsA • PsO • RA • UC • UV	SC injection: • Self-administered	SC: • 40 mg/0.8 mL PEN • 40 mg/0.8 mL PFS	• Refrigerate at 36°F to 46°F (2°C to 8°C) • Do not freeze • Store in original packages • Protect from light	Infections (e.g., upper respiratory, sinusitis), injection site reactions, headache and rash

Autoimmune (AI) Drug List

Select FDA-approved AI specialty medications and infusion therapy

Drug name	Manufacturer	FDA-Approved Indication(s)	Route of Administration	Supply Method	Storage	Select Common Adverse Reactions
Adalimumab-aaty (unbranded version of Yuflyma)	Celltrion USA	SC injection: • AS • CD (6+) • HS • PJIA (2+) • PsA • PsO • RA • UC • UV	SC injection: • Self-administered	SC: • 20 mg/0.2 mL PFS • 40 mg/0.4 mL PFS • 40 mg/0.4 mL PEN • 80 mg/0.8 mL PEN	• Refrigerate at 36°F to 46°F (2°C to 8°C) • Do not freeze • Store in original packages • Protect from light	Infections (e.g., upper respiratory, sinusitis), injection site reactions, headache and rash
Adalimumab-adaz (unbranded version of Hyrimoz)	Sandoz	• AS • CD (6+) • HS • PJIA (2+) • PsO • PsA • RA • UC • UV	SC injection: • Self-administered	SC: • 10 mg/0.1 mL PFS • 20 mg/0.2 mL PFS • 40 mg/0.4 mL PFS • 40 mg/0.4 mL PEN • 80 mg/0.8 mL PEN	• Refrigerate at 36°F to 46°F (2°C to 8°C) • Do not freeze • Store in original packages • Protect from light	Infections (e.g., upper respiratory, sinusitis), injection site reactions, headache and rash

Autoimmune (AI) Drug List

Select FDA-approved AI specialty medications and infusion therapy

Drug name	Manufacturer	FDA-Approved Indication(s)	Route of Administration	Supply Method	Storage	Select Common Adverse Reactions
Adalimumab-fkjp (unbranded version of Huloio)	Biocon	• AS • CD (6+) • HS • PJIA (2+) • PsO • PsA • RA • UC • UV	SC injection: • Self-administered	SC: • 40 mg/0.8 mL PEN • 40 mg/0.8 mL PFS • 20 mg/0.4 mL PFS	• Refrigerate at 36°F to 46°F (2°C to 8°C) • Do not freeze • Protect from light • Store in original package	Infections (e.g., upper respiratory, sinusitis), injection site reactions, headache and rash
Adbry (tralokinumab-ldrm)	LEO Pharma	• AD (12+)	SC injection: • Self-administered	SC: • 150 mg/mL PFS • 300 mg/2 mL PEN	• Store refrigerated at 36°F to 46°F (2°C to 8°C) • Store in the original carton to protect from light	Upper respiratory tract infections, conjunctivitis, injection site reactions and eosinophilia
Amjevita (adalimumab-atto)	Amgen	• AS • CD (6+) • HS • PJIA (2+) • PsO • PsA • RA • UC • UV	SC injection: • Self-administered	SC: • 80 mg/0.8 mL PEN • 40 mg/0.4 mL PEN • 40 mg/0.4 mL PFS • 20 mg/0.2 mL PFS • 10 mg/0.2 mL PFS	• Refrigerate at 36°F to 46°F (2°C to 8°C) • Do not freeze • Protect from light • Store in original package	Infections (e.g., upper respiratory, sinusitis), injection site reactions, headache and rash

Autoimmune (AI) Drug List

Select FDA-approved AI specialty medications and infusion therapy

Drug name	Manufacturer	FDA-Approved Indication(s)	Route of Administration	Supply Method	Storage	Select Common Adverse Reactions
Arcalyst (rilonacept)	Kiniksa Pharmaceuticals	• CAPS (12+) • FCAS (12+) • MWS (12+) • DIRA (≥ 10 kg) • RP (12+)	SC injection: • Self-administered	SC: • 220 mg of lyophilized powder in a single dose vial for reconstitution	• Refrigerate at 36°F to 46°F (2°C to 8°C) • Store in original carton • Protect from light • Do not use beyond the date stamped on the label • After reconstitution, may be kept at room temperature, protected from light	Injection-site reactions and upper respiratory tract infections
Avsola (infliximab-axxq)	Amgen	• AS • CD (6+) • PsO • PsA • RA • UC (6+)	IV infusion: • Prepared and infused by a health care professional	IV: • 100 mg lyophilized powder in a single dose vial for reconstitution	• Store unopened Avsola vials in a refrigerator at 36°F to 46°F (2°C to 8°C) • Protect from light	Infections (e.g., upper respiratory, sinusitis and pharyngitis), infusion-related reactions, headache and abdominal pain

Autoimmune (AI) Drug List

Select FDA-approved AI specialty medications and infusion therapy

Drug name	Manufacturer	FDA-Approved Indication(s)	Route of Administration	Supply Method	Storage	Select Common Adverse Reactions
Benlysta (belimumab)	GlaxoSmithKline	• SLE (5+) • LN (5+)	IV infusion: • Prepared and infused by a health care professional SC injection: • Self-administered	IV: • 120 mg lyophilized powder in a single dose vial for reconstitution • 400 mg lyophilized powder in a single dose vial for reconstitution SC: • 200 mg/mL PEN • 200 mg/mL PFS	• Refrigerate at 36°F to 46°F (2°C to 8°C) • Do not freeze • Store in original packages • Protect from light • Avoid exposure to heat	Nausea, diarrhea, pyrexia, nasopharyngitis, bronchitis, pharyngitis, insomnia, pain in extremity, depression and migraine
Bimzelx (bimekizumab-bkzx)	UCB	• AS • HS • Nr-axSPA • PsO • PsA	SC injection: • Self-administered	SC: • 160mg/mL PEN • 160mg/mL PFS • 320 mg/2 mL PEN • 320 mg/2 mL PFS	• Refrigerate at 36°F to 46°F (2°C to 8°C) • Do not freeze • Do not shake • Store in original carton to protect from light	Upper respiratory tract infections, oral candidiasis, headache, injection site reactions, tinea infections, gastroenteritis, Herpes simplex infections, acne, folliculitis, other candida infections and fatigue

Autoimmune (AI) Drug List

Select FDA-approved AI specialty medications and infusion therapy

Drug name	Manufacturer	FDA-Approved Indication(s)	Route of Administration	Supply Method	Storage	Select Common Adverse Reactions
Cibinqo (abrocitinib)	Pfizer	AD (12+)	Oral	Oral: 50 mg tablets 100 mg tablets 200 mg tablets	- Store at 68°F to 77°F (20°C to 25°C) - Keep in original package	Nasopharyngitis, nausea, headache, herpes simplex, increased blood creatinine phosphokinase, dizziness, urinary tract infection, fatigue, acne, vomiting, oropharyngeal pain, influenza, gastroenteritis, impetigo, hypertension, contact dermatitis, upper abdominal pain, abdominal discomfort, herpes zoster and thrombocytopenia
Cimzia (certolizumab pegol)	UCB	- AS - CD - Nr-axSpA - PJIA (2+) - PsO - PsA - RA	SC injection: Self-administered (PFS only)	SC: 200 mg/mL PFS 200 mg lyophilized powder in a single-dose vial (for health care professional use only)	- Refrigerate at 36°F to 46°F (2°C to 8°C) - Do not freeze - Protect from light	Upper respiratory tract infection, rash and urinary tract infection
Cosentyx (secukinumab)	Novartis	SC: - AS - ERA (4+) - HS - Nr-axSpA - PsO (6+) - PsA (2+)	SC injection: Self-administered	SC: 75 mg/0.5 mL PFS 150 mg/mL PEN 150 mg/mL PFS 300 mg/2 mL PEN 300 mg/2 mL PFS	SC: - Refrigerate at 36°F to 46°F (2°C to 8°C) - Do not freeze - Keep the product in the original carton - Protect from light - Do not shake to avoid foaming	Nasopharyngitis, diarrhea and upper respiratory tract infection

Autoimmune (AI) Drug List

Select FDA-approved AI specialty medications and infusion therapy

Drug name	Manufacturer	FDA-Approved Indication(s)	Route of Administration	Supply Method	Storage	Select Common Adverse Reactions
Dupixent (dupilumab)	Sanofi	• AD (6 mo+) • Asthma (6+) • COPD • CRSwNP (12+) • CRSwNP • EoE (1+ and ≥ 15 kg) • PN	SC injection: • Self-administered	SC: • 200 mg/1.14 mL PEN • 200 mg/1.14 mL PFS • 300 mg/2 mL PEN • 300 mg/2 mL PFS	• Refrigerate at 36°F to 46°F (2°C to 8°C) in the original carton to protect from light • Do not freeze	Injection site reactions, conjunctivitis, blepharitis, oral herpes, keratitis, eye pruritus, other herpes simplex virus infection and dry eye
Ebglyss (lebrikizumab-lbkz)	Eli Lilly	• AD (12+)	SC injection: • Self-administered	SC: • 250 mg/2 mL PFS • 250 mg/2 mL PEN	• Refrigerate at 36°F to 46°F (2°C to 8°C) • Store in the original carton to protect from light • Do not freeze • Do not shake	Conjunctivitis, injection site reactions, and herpes zoster
Enbrel (etanercept)	Amgen	• AS • JPsA (2+) • PsO (4+) • PsA (2+) • RA	SC injection: • Self-administered	SC: • 25 mg/0.5 mL single dose vial • 25 mg/0.5 mL PFS • 50 mg/mL PFS • 50 mg/mL PEN • 50 mg/mL Enbrel Mini single-dose prefilled cartridge	• Refrigerate at 36°F to 46°F (2°C to 8°C) • Do not freeze • Do not shake • Store in the original carton to protect from light or physical damage	Infections and injection site reactions

Autoimmune (AI) Drug List

Select FDA-approved AI specialty medications and infusion therapy

Drug name	Manufacturer	FDA-Approved Indication(s)	Route of Administration	Supply Method	Storage	Select Common Adverse Reactions
Entyvio (vedolizumab)	Takeda	• CD • UC	IV infusion: • HCP-administered SC injection: • Self-administered	IV: • 300 mg single dose vial SC: • 108 mg/0.68 mL PEN	• Refrigerate unopened vials, PEN, and PFS at 36°F to 46°F (2°C to 8°C) • Do not freeze • Retain in original package to protect from light	Nasopharyngitis, headache, arthralgia, nausea, pyrexia, upper respiratory tract infection, fatigue, cough, bronchitis, influenza, back pain, rash, pruritus, sinusitis, oropharyngeal pain and pain in extremities. In addition, injection site reactions are associated with the SC injection only.
Hadlima (adalimumab-bwwd)	Samsung Bioepis/ Organon	• AS • CD (6+) • HS • PJIA (2+) • PsO • PsA • RA • UC • UV	SC injection: • Self-administered	SC: • 40 mg/0.8 mL PFS • 40 mg/0.4 mL PFS • 40 mg/0.8 mL PEN • 40 mg/0.4 mL PEN	• Refrigerate at 36°F to 46°F (2°C to 8°C) • Do not freeze • Protect from light • Store in original container	Infections (e.g., upper respiratory, sinusitis), injection site reactions, headache and rash
Hulio (adalimumab-fkjp)	Mylan	• AS • CD (6+) • HS • PJIA (2+) • PsO • PsA • RA • UC • UV	SC injection: • Self-administered	SC: • 40 mg/0.8 mL PEN • 40 mg/0.8 mL PFS • 20 mg/0.4 mL PFS	• Refrigerate at 36°F to 46°F (2°C to 8°C) • Do not freeze • Protect from light • Store in original carton	Infections (e.g., upper respiratory, sinusitis), injection site reactions, headache and rash

Autoimmune (AI) Drug List

Select FDA-approved AI specialty medications and infusion therapy

Drug name	Manufacturer	FDA-Approved Indication(s)	Route of Administration	Supply Method	Storage	Select Common Adverse Reactions
Humira (adalimumab)	AbbVie/ Cordavis	• AS • CD (6+) • HS (12+) • PJIA (2+) • PsO • PsA • RA • UC (5+) • UV (2+)	SC injection: • Self-administered	SC: • 80 mg/0.8 mL PEN • 40 mg/0.8 mL PEN • 40 mg/0.4 mL PEN • 40 mg/0.8 mL PFS • 40 mg/0.4 mL PFS • 20 mg/0.2 mL PFS • 10 mg/0.1 mL PFS	• Refrigerate at 36°F to 46°F (2°C to 8°C) • Do not freeze • Protect from light • Store in original carton	Infections (e.g., upper respiratory, sinusitis), injection site reactions, headache and rash
Hyrimoz (adalimumab-adaz)	Cordavis/ Sandoz	• AS • CD (6+) • HS • PJIA (2+) • PsO • PsA • RA • UC • UV	SC injection: • Self-administered	SC: • 80 mg/0.8 mL PEN • 40 mg/0.4 mL PEN • 40 mg/0.8 mL PEN • 40 mg/0.4 mL PFS • 40 mg/0.8 mL PFS • 20 mg/0.2 mL PFS	• Refrigerate at 36°F to 46°F (2°C to 8°C) • Do not freeze • Protect from light • Store in original carton	Infections (e.g., upper respiratory, sinusitis), injection site reactions, headache and rash

Autoimmune (AI) Drug List

Select FDA-approved AI specialty medications and infusion therapy

Drug name	Manufacturer	FDA-Approved Indication(s)	Route of Administration	Supply Method	Storage	Select Common Adverse Reactions
Ilaris (canakinumab)	Novartis	• CAPS (4+) • FCAS (4+) • FMF • Gout flares • HIDS/MKD • MWS (4+) • TRAPS • Active Still's Disease/AOSD (2+) • SJIA (2+)	SC injection: • HCP-administered	SC: • 150 mg/mL single dose vial	• Refrigerate at 36°F to 46°F (2°C to 8°C) • Do not freeze • Store in the original carton to protect from light	CAPS: Nasopharyngitis, diarrhea, influenza, rhinitis, nausea, headache, bronchitis, gastroenteritis, pharyngitis, weight increased, musculoskeletal pain and vertigo TRAPS, HIDS/MKD, and FMF: injection-site reactions and nasopharyngitis Still's Disease: Nasopharyngitis and upper respiratory tract infections, abdominal pain and injection-site reactions Gout Flares: Nasopharyngitis, upper respiratory tract infections, urinary tract infections, hypertriglyceridemia and back pain
Ilumya (tildrakizumab-asmn)	Sun Pharma	PsO	SC injection: • HCP-administered	SC: • 100 mg/mL PFS	• Refrigerate at 36°F to 46°F (2°C to 8°C) in the original carton to protect from light until the time of use • Do not freeze • Do not shake	Upper respiratory infections, injection site reactions and diarrhea

Autoimmune (AI) Drug List

Select FDA-approved AI specialty medications and infusion therapy

Drug name	Manufacturer	FDA-Approved Indication(s)	Route of Administration	Supply Method	Storage	Select Common Adverse Reactions
Inflectra (infliximab-dyyb)	Celltrion/Pfizer	- AS - CD (6+) - PsO - PsA - RA - UC (6+)	IV infusion: Prepared and infused by a health care professional	IV: 100 mg lyophilized in a powder single dose vial for reconstitution	Refrigerate at 36°F to 46°F (2°C to 8°C)	Infections (e.g., upper respiratory, sinusitis and pharyngitis), infusion-related reactions, headache and abdominal pain
Infliximab (unbranded version of Remicade)	Janssen	- AS - CD (6+) - PsO - PsA - RA - UC (6+)	IV infusion: Prepared and infused by a health care professional	IV: 100 mg lyophilized in a powder single dose vial for reconstitution	Refrigerate at 36°F to 46°F (2°C to 8°C)	Infections (e.g., upper respiratory, sinusitis and pharyngitis), infusion-related reactions, headache and abdominal pain
Kevzara (sarilumab)	Sanofi	- RA - PJIA (> 63 kg) - PMR	SC injection: Self-administered	SC: 150 mg/1.14 mL PFS 200 mg/1.14 mL PFS 150 mg/1.14 mL PEN 200 mg/1.14 mL PEN	- Refrigerate at 36°F to 46°F (2°C to 8°C) in original carton to protect from light - Do not freeze - Do not shake	PJIA: nasopharyngitis, neutropenia, upper respiratory tract infection and injection site erythema PMR: Neutropenia, leukopenia and injection site pruritus RA: Neutropenia, increased ALT, injection site erythema, upper respiratory infections and urinary tract infections

Autoimmune (AI) Drug List

Select FDA-approved AI specialty medications and infusion therapy

Drug name	Manufacturer	FDA-Approved Indication(s)	Route of Administration	Supply Method	Storage	Select Common Adverse Reactions
Leqselvi (deuruxolitinib)	Sun Pharma	• AA	Oral	Oral: • 8 mg tablets	• Store at 68°F to 77°F (20°C to 25°C) • Store in original package	Acne, headache, nasopharyngitis, blood creatine phosphokinase increased, hyperlipidemia, fatigue, weight increased, lymphopenia, thrombocytosis, anemia, skin and soft tissue infections, neutropenia, and herpes
Litfulo (ritlecitinib)	Pfizer	• AA (12+)	Oral	Oral: • 50 mg capsules	• Store at 68°F to 77°F (20°C to 25°C) • Store in original package	Headache, diarrhea, acne, rash, urticaria, folliculitis, pyrexia, atopic dermatitis, dizziness, blood creatine phosphokinase increased, herpes zoster, red blood cell count decreased and stomatitis
Nemluvio (nemolizumab-ilto)	Galderma	• AD (12+) • PN	SC injection: • Self-administered	SC: • 30 mg/0.49 mL PEN	• Refrigerate at 36°F to 46°F (2°C to 8°C) in original carton to protect from light • Do not freeze • Do not expose to heat or direct sunlight	AD: Headache (including migraine), arthralgia, urticaria, and myalgia PN: Headache, dermatitis, atopic eczema, and eczema nummular

Autoimmune (AI) Drug List

Select FDA-approved AI specialty medications and infusion therapy

Drug name	Manufacturer	FDA-Approved Indication(s)	Route of Administration	Supply Method	Storage	Select Common Adverse Reactions
Olumiant (baricitinib)	Eli Lilly	• AA • RA • Hospitalized adults who have COVID-19	Oral	Oral: • 1 mg tablets • 2 mg tablets • 4 mg tablets	• Store at 68°F to 77°F (20°C to 25°C)	RA: upper respiratory tract infections, nausea, herpes simplex and herpes zoster COVID-19: increases of liver enzymes, thrombocytosis, creatine phosphokinase increases, neutropenia, deep vein thrombosis, pulmonary embolism and urinary tract infection AA: upper respiratory tract infections, headache, acne, hyperlipidemia, creatine phosphokinase increase, urinary tract infection, liver enzyme elevations, folliculitis, fatigue, lower respiratory tract infections, nausea, genital Candida infections, anemia, neutropenia, abdominal pain, herpes zoster and weight increase

Autoimmune (AI) Drug List

Select FDA-approved AI specialty medications and infusion therapy

Drug name	Manufacturer	FDA-Approved Indication(s)	Route of Administration	Supply Method	Storage	Select Common Adverse Reactions
Omvo (mirikizumab-mrkz)	Eli Lilly	- CD - UC	IV infusion: - HCP-administered SC injection: - Self-administered	IV: 300 mg/15 mL single dose vial SC: 100 mg/mL PEN 100 mg/mL PFS	- Refrigerate at 36°F to 46°F (2°C to 8°C) - Store in original container to protect from light - Do not shake - Do not freeze	CD: Upper respiratory tract infections, injection site reactions, headache, arthralgia, and elevated liver tests UC Induction: Upper respiratory tract infections and arthralgia. UC Maintenance: Upper respiratory tract infections, injection site reactions, arthralgia, rash, headache and herpes viral infection
Orencia (abatacept)	Bristol Myers Squibb	IV: - PJIA (6+) - Prophylaxis of aGVHD (2+) - PsA - RA SC: - PJIA (2+) - PsA (2+) - RA	IV infusion: - Prepared and infused by a health care professional SC injection: - Self-administered	IV: 250 mg lyophilized powder in a single dose vial for reconstitution SC: 50 mg/0.4 mL PFS 87.5 mg/0.7 mL PFS 125 mg/mL PFS 125 mg/mL PEN	IV: - Refrigerate at 36°F to 46°F (2°C to 8°C) - Store in original container - Protect from light SC: - Refrigerate at 36°F to 46°F (2°C to 8°C) - Store in original container - Protect from light - Do not freeze	IV: RA: headache, upper respiratory tract infection, nasopharyngitis and nausea aGVHD: anemia, hypertension, CMV reactivation/CMV infection, pyrexia, pneumonia, epistaxis, CD4 lymphocytes decreased, hypermagnesemia and acute kidney injury SC: Headache, upper respiratory tract infection, nasopharyngitis and nausea

Autoimmune (AI) Drug List

Select FDA-approved AI specialty medications and infusion therapy

Drug name	Manufacturer	FDA-Approved Indication(s)	Route of Administration	Supply Method	Storage	Select Common Adverse Reactions
Otezla (apremilast)	Amgen	• Oral ulcers associated with Behçet's Disease • PsA • PsO (6+)	Oral	Oral: • 28 day Starter Pack: 10 mg, 20 mg, and 30 mg tablets • 20 mg tablets • 30 mg tablets	• Store below 86°F (30°C)	PsA: diarrhea, nausea and headache PsO: diarrhea, nausea, upper respiratory tract infection and headache, including tension headache Behçet's Disease: diarrhea, nausea, headache and upper respiratory tract infection
Otrexup (methotrexate)	Assertio Pharmaceuticals	• PJIA • PsO • RA	SC injection: • Self-administered	SC: • 10 mg/0.4 mL PEN • 12.5 mg/0.4 mL PEN • 15 mg/0.4 mL PEN • 17.5 mg/0.4 mL PEN • 20 mg/0.4 mL PEN • 22.5 mg/0.4 mL PEN • 25 mg/0.4mL PEN	• Store at controlled room temperature, 77°F (25°C); excursions permitted to 59°F to 86°F (15°C to 30°C) • Protect from light	Nausea, abdominal pain, dyspepsia, stomatitis/mouth sores, rash, nasopharyngitis, diarrhea, liver function test abnormalities, vomiting, headache, bronchitis, thrombocytopenia, alopecia, leucopenia, pancytopenia, dizziness, photosensitivity and "burning of skin lesions"

Autoimmune (AI) Drug List

Select FDA-approved AI specialty medications and infusion therapy

Drug name	Manufacturer	FDA-Approved Indication(s)	Route of Administration	Supply Method	Storage	Select Common Adverse Reactions
Pyzchiva (ustekinumab-ttwe)	Samsung Bioepis	• CD • PsA (6+) • PsO (6+) • UC	IV infusion: • HCP-administered SC injection: • Self-administered	IV: • 130 mg/26 mL SDV SC: • 45mg/0.5 mL PFS • 90 mg/mL PFS	• Refrigerate 36°F to 46°F (2°C to 8°C) • Do not freeze • Do not shake • Store in original carton to protect from light	CD Induction: Vomiting CD Maintenance: Nasopharyngitis, injection, site erythema, vulvovaginal, candidiasis/ mycotic, infection, bronchitis, pruritus, urinary tract, and sinusitis PsO: Nasopharyngitis, upper respiratory tract infection, headache and fatigue UC Induction: Nasopharyngitis UC Maintenance: Nasopharyngitis, headache, abdominal pain, influenza, fever, diarrhea, sinusitis, fatigue and nausea

Autoimmune (AI) Drug List

Select FDA-approved AI specialty medications and infusion therapy

Drug name	Manufacturer	FDA-Approved Indication(s)	Route of Administration	Supply Method	Storage	Select Common Adverse Reactions
Rasuvo (methotrexate)	Medexus	• PJIA • PsO • RA	SC injection: • Self-administered	SC: • 7.5 mg/0.15 mL PEN • 10 mg/0.20 mL PEN • 12.5 mg/0.25 mL PEN • 15 mg/0.30 mL PEN • 17.5 mg/0.35 mL PEN • 20 mg/0.40 mL PEN • 22.5mg/0.45mL PEN • 25 mg/0.50 mL PEN • 30 mg/0.60 mL PEN	• Store at controlled room temperature, 77°F (25°C) • Protect from light	Nausea, abdominal pain, dyspepsia, stomatitis/mouth sores, rash, nasopharyngitis, diarrhea, liver function test abnormalities, vomiting, headache, bronchitis, thrombocytopenia, alopecia, leucopenia, pancytopenia, dizziness, photosensitivity and “burning of skin lesions”
Remicade (infliximab)	Janssen	• AS • CD (6+) • PsO • PsA • RA • UC (6+)	IV infusion: • Prepared and infused by a health care professional	IV: • 100 mg lyophilized powder in a single dose vial for reconstitution	• Refrigerate at 36°F to 46°F (2°C to 8°C)	Infections (e.g., upper respiratory, sinusitis and pharyngitis), infusion-related reactions, headache and abdominal pain

Autoimmune (AI) Drug List

Select FDA-approved AI specialty medications and infusion therapy

Drug name	Manufacturer	FDA-Approved Indication(s)	Route of Administration	Supply Method	Storage	Select Common Adverse Reactions
Renflexis (infliximab-abda)	Samsung Bioepis/ Organon	• AS • CD (6+) • PsO • PsA • RA • UC (6+)	IV infusion: • Prepared and infused by a health care professional	IV: • 100 mg lyophilized powder in a single dose vial for reconstitution	• Refrigerate at 36°F to 46°F (2°C to 8°C)	Infections (e.g., upper respiratory, sinusitis and pharyngitis), infusion-related reactions, headache and abdominal pain
Riabni (rituximab-arrx)	Amgen	• RA	IV infusion only: • HCP-administered • Do not administer by IV push or bolus	IV: • 100 mg/10 mL single dose vial • 500 mg/50 mL single dose vial	• Refrigerate vials at 36°F to 46°F (2°C to 8°C) • Do not freeze or shake • Protect from direct sunlight	Upper respiratory tract infection, nasopharyngitis, urinary tract infection and bronchitis (other important adverse reactions include infusion-related reactions, serious infections and cardiovascular events)

Autoimmune (AI) Drug List

Select FDA-approved AI specialty medications and infusion therapy

Drug name	Manufacturer	FDA-Approved Indication(s)	Route of Administration	Supply Method	Storage	Select Common Adverse Reactions
Rinvoq (upadacitinib)	AbbVie	Rinvoq: • AD (12+) • AS • CD • GCA • Nr-axSpA • PJIA (2+) • PsA (2+) • RA • UC Rinvoq LQ: • PJIA (2+) • PsA (2+)	Oral	Oral: • Extended-release tablets: 15 mg, 30 mg, 45 mg Oral Solution (LQ): • 1 mg/1 mL oral solution	• Store at 36°F to 77°F (2°C to 25°C) • Store in the original bottle to protect from moisture (oral tablet only)	AD: Upper respiratory tract infections, acne, herpes simplex, headache, blood creatine phosphokinase increased, cough, hypersensitivity, folliculitis, nausea, abdominal pain, pyrexia, increased weight, herpes zoster, influenza, fatigue, neutropenia, myalgia and influenza like illness CD: Upper respiratory tract infections, anemia, pyrexia, acne, herpes zoster and headache GCA: Upper respiratory tract infections, headache, fatigue, peripheral edema, cough, anemia, rash, herpes zoster, and nausea UC: Upper respiratory tract infections, increased blood creatine phosphokinase, acne, neutropenia, elevated liver enzymes, pyrexia, and rash RA/PsA/AS/nr-axSpA: Upper respiratory tract infections, herpes zoster, herpes simplex, bronchitis, nausea, cough, pyrexia, acne and headache

Autoimmune (AI) Drug List

Select FDA-approved AI specialty medications and infusion therapy

Drug name	Manufacturer	FDA-Approved Indication(s)	Route of Administration	Supply Method	Storage	Select Common Adverse Reactions
Rituxan (rituximab)	Genentech	RA	IV infusion: HCP-administered	IV: 100 mg/10 mL single dose vial 500 mg/50 mL single dose vial	- Refrigerate vials at 36°F to 46°F (2°C to 8°C) - Do not freeze or shake - Protect from direct sunlight	Upper respiratory tract infection, nasopharyngitis, urinary tract infection and bronchitis (other important adverse reactions include infusion-related reactions, serious infections and cardiovascular events)
Ruxience (rituximab-pvvr)	Pfizer	RA	IV infusion: HCP-administered	IV: 100 mg/10 mL single dose vial 500 mg/50 mL single dose vial	- Refrigerate vials at 36°F to 46°F (2°C to 8°C) - Do not freeze or shake - Protect from direct sunlight	Upper respiratory tract infection, nasopharyngitis, urinary tract infection and bronchitis (other important adverse reactions include infusion-related reactions, serious infections and cardiovascular events)
Saphnelo (anifrolumab-fnia)	AstraZeneca	SLE	IV Infusion: HCP-administered	IV: 300 mg/2 mL single dose vial	- Store in a refrigerator at 36°F to 46°F (2°C to 8°C) - Store in the original carton to protect from light - Do not freeze - Do not shake	Nasopharyngitis, upper respiratory tract infections, bronchitis, infusion related reactions, herpes zoster and cough

Autoimmune (AI) Drug List

Select FDA-approved AI specialty medications and infusion therapy

Drug name	Manufacturer	FDA-Approved Indication(s)	Route of Administration	Supply Method	Storage	Select Common Adverse Reactions
Selarsdi (ustekinumab-aekn)	Alvotech	• CD • PsA (6+) • PsO (6+) • UC	IV infusion: • HCP-administered SC injection: • Self-administered	IV: • 130 mg/26 mg SDV SC: • 45 mg/0.5 mL PFS • 90 mg/mL PFS	• Refrigerate 36°F to 46°F (2°C to 8°C) • Do not freeze • Do not shake • Store in original carton to protect from light	CD Induction: Vomiting CD Maintenance: Nasopharyngitis, injection site erythema, vulvovaginal candidiasis/ mycotic, infection, bronchitis, pruritus, urinary tract infection and sinusitis PsO: Nasopharyngitis, upper respiratory tract infection, headache and fatigue UC Induction: Nasopharyngitis UC Maintenance: Nasopharyngitis, headache, abdominal pain, influenza, fever, diarrhea, sinusitis, fatigue and nausea
Siliq (brodalumab)	Valeant/Ortho Dermatologics	• PsO	SC injection: • Self-administered	SC injection: • Self-administered	• Store refrigerated at 36°F to 46°F (2°C to 8°C) • Store in the original carton to protect from light • Do not freeze • Do not shake	Arthralgia, headache, fatigue, diarrhea, oropharyngeal pain, nausea, myalgia, injection site reactions, influenza, neutropenia and tinea infections

Autoimmune (AI) Drug List

Select FDA-approved AI specialty medications and infusion therapy

Drug name	Manufacturer	FDA-Approved Indication(s)	Route of Administration	Supply Method	Storage	Select Common Adverse Reactions
Simlandi (adalimumab-ryvk)	Alvotect/Teva	• AS • CD (6+) • HS • PJIA (2+) • PsO • PsA • RA • UC • UV	SC injection: • Self-administered	SC: • 20 mg/0.2 mL PFS • 40 mg/0.4 mL PEN • 40 mg/0.4 mL PFS • 80 mg/0.4 mL PEN	• Refrigerate at 36°F to 46°F (2°C to 8°C) • Do not freeze • Protect from light • Store in original carton	Infections (e.g., upper respiratory, sinusitis), injection site reactions, headache and rash
Simponi ARIA (golimumab)	Janssen	• AS • PJIA (2+) • PsA (2+) • RA	IV infusion only: • HCP-administered	IV: • 50 mg/4 mL single dose vial	• Refrigerate at 36°F to 46°F (2°C to 8°C) • Do not freeze • Do not shake • Store in original carton to protect from light	Upper respiratory tract infection, alanine aminotransferase increased, viral infection, aspartate aminotransferase increased, neutrophil count decreased, bronchitis, hypertension and rash

Autoimmune (AI) Drug List

Select FDA-approved AI specialty medications and infusion therapy

Drug name	Manufacturer	FDA-Approved Indication(s)	Route of Administration	Supply Method	Storage	Select Common Adverse Reactions
Skyrizi (risankizumab-rzaa)	AbbVie	IV/OBI: - CD - UC SC: - PsA - PsO	IV infusion: - HCP-administered SC injection: - Self-administered	IV: 600 mg/10 mL single dose vial SC: 360 mg/2.4 mL OBI 180 mg/1.2 mL OBI 150 mg/mL PEN 150 mg/mL PFS	- Store in a refrigerator at 36°F to 46° F (2°C to 8°C) - Do not freeze - Do not shake - Store in the original cartons to protect from light	CD Induction: Upper respiratory infections, headache, and arthralgia CD Maintenance: Arthralgia, abdominal pain, injection site reactions, anemia, pyrexia, back pain, arthropathy and urinary tract infection PsO and PsA: Upper respiratory infections, headache, fatigue, injection site reactions and tinea infections UC Induction: Arthralgia UC Maintenance: Arthralgia, pyrexia, injection site reactions, and rash
Sotyktu (deucravacitinib)	Bristol Meyers Squibb	PsO	Oral	Oral: 6 mg tablets	Store at 68°F to 77°F (20°C to 25°C)	Upper respiratory infections, blood creatine phosphokinase increased, herpes simplex, mouth ulcers, folliculitis and acne

Autoimmune (AI) Drug List

Select FDA-approved AI specialty medications and infusion therapy

Drug name	Manufacturer	FDA-Approved Indication(s)	Route of Administration	Supply Method	Storage	Select Common Adverse Reactions
Stelara (ustekinumab)	Janssen	• CD • PsA (6+) • PsO (6+) • UC	IV infusion: • Prepared and infused by a health care professional SC injection: • Self-administered	IV: • 130 mg/26 mL SDV SC: • 45 mg/0.5 mL PFS • 90 mg/mL PFS • 45 mg/0.5 mL SDV	• Refrigerate 36°F to 46°F (2°C to 8°C) • Do not freeze • Do not shake • Store in original carton to protect from light	PsO and PsA: Nasopharyngitis, upper respiratory tract infection, headache and fatigue CD Induction: Vomiting CD Maintenance: Nasopharyngitis, injection site erythema, vulvovaginal candidiasis/ mycotic infection, bronchitis, pruritus, urinary tract infection, and sinusitis UC Induction: Nasopharyngitis UD Maintenance: Nasopharyngitis, headache, abdominal pain, influenza, fever, diarrhea, sinusitis, fatigue and nausea

Autoimmune (AI) Drug List

Select FDA-approved AI specialty medications and infusion therapy

Drug name	Manufacturer	FDA-Approved Indication(s)	Route of Administration	Supply Method	Storage	Select Common Adverse Reactions
Steqeyma (ustekinumab-stba)	Celltrion	• CD • PsA (6+) • PsO (6+) • UC	IV infusion: • HCP-administered SC injection: • Self-administered	IV: • 130 mg/26 mL SDV SC: • 45 mg/0.5 mL PFS • 90 mg/mL PFS	• Refrigerate 36°F to 46°F (2°C to 8°C) • Do not freeze • Do not shake • Store in original carton to protect from light	CD Induction: Vomiting CD Maintenance: Nasopharyngitis, injection site erythema, vulvovaginal candidiasis/ mycotic infection, bronchitis, pruritus, urinary tract infection and sinusitis PsO: Nasopharyngitis, upper respiratory tract infection, headache and fatigue UC Induction: Nasopharyngitis. CD Maintenance: Nasopharyngitis, headache, abdominal pain, influenza, fever, diarrhea, sinusitis, fatigue, and nausea
Taltz (ixekizumab)	Eli Lilly	• AS • Nr-axSpA • PsA • PsO (6+)	SC injection: • Self-administered	SC: • 20 mg/0.25 mL PFS • 40 mg/0.5 mL PFS • 80 mg/mL PEN • 80 mg/mL PFS	• Refrigerate 36°F to 46°F (2°C to 8°C) • Do not freeze • Protect from light • Do not shake	Injection site reactions, upper respiratory tract infections, nausea and tinea infections

Autoimmune (AI) Drug List

Select FDA-approved AI specialty medications and infusion therapy

Drug name	Manufacturer	FDA-Approved Indication(s)	Route of Administration	Supply Method	Storage	Select Common Adverse Reactions
Tremfya (guselkumab)	Janssen	• CD • PsA • PsO • UC	SC injection: • Self-administered IV infusion: • Prepared and infused by a health care professional	SC: • 100 mg/mL PFS • 100 mg/mL PEN • 200 mg/2 mL PFS • 200 mg/2 mL PEN IV: • 200 mg/20 mL single dose vial	• Refrigerate at 36°F to 46°F (2°C to 8°C) • Store in original carton to protect from light • Do not freeze • Do not shake	Upper respiratory infections, headache, injection site reactions, arthralgia, bronchitis, diarrhea, gastroenteritis, tinea infections and herpes simplex infections
Truxima (rituximab-abbs)	Celltrion/Teva	• RA	IV infusion: • HCP-administered	IV: • 100 mg/10 mL single dose vial • 500 mg/50 mL single dose vial	• Refrigerate vials at 36°F to 46°F (2°C to 8°C) • Do not freeze or shake • Protect from direct sunlight	Upper respiratory tract infection, nasopharyngitis, urinary tract infection and bronchitis (other important adverse reactions include infusion-related reactions, serious infections and cardiovascular events)
Tyenne (tocilizumab-aazg)	Fresenius Kabi	• GCA • PJIA (2+) • RA • SJIA (2+)	SC Injection: • Self-administered IV infusion: • Prepared and infused by a health care professional	SC: • 162 mg/0.9 mL PFS • 162 mg/0.9 mL PEN IV: • 80 mg/4 mL single dose vial • 200 mg/10 mL single dose vial • 400 mg/20 mL single dose vial	• Refrigerate at 36°F to 46°F (2°C to 8°C) • Do not freeze • Store in original packages • Protect from light • Keep syringes and pen dry	Upper respiratory tract infections, nasopharyngitis, headache, hypertension, increased alanine aminotransferase (ALT) and injection site reactions

Autoimmune (AI) Drug List

Select FDA-approved AI specialty medications and infusion therapy

Drug name	Manufacturer	FDA-Approved Indication(s)	Route of Administration	Supply Method	Storage	Select Common Adverse Reactions
Velsipity (etrasimod)	Pfizer	• UC	Oral	Oral: • 2 mg tablet	• Store at room temperature at 68°F to 77°F (20°C to 25°C); excursions permitted between 59°F to 86°F (15°C to 30°C)	Headache, elevated liver tests and dizziness
Xeljanz/Xeljanz XR (tofacitinib)	Pfizer	• AS • PJIA (2+) • PsA • RA • UC	Oral	Oral: • Tablets: – 5 mg – 10 mg • XR tablets: – 11 mg – 22 mg • Oral Solution: – 1 mg/mL	• Store at room temperature at 68°F to 77°F (20°C to 25°C) • Do not repackage (oral tablet only) • Store in original package and protect from light (oral solution only)	RA, PsA, AS and PJIA: Upper respiratory tract infection, nasopharyngitis, diarrhea and headache UC: Nasopharyngitis, elevated cholesterol levels, headache, upper respiratory tract infection, increased blood creatine phosphokinase, rash, diarrhea and herpes zoster

Autoimmune (AI) Drug List

Select FDA-approved AI specialty medications and infusion therapy

Drug name	Manufacturer	FDA-Approved Indication(s)	Route of Administration	Supply Method	Storage	Select Common Adverse Reactions
Yesintek (ustekinumab-kfce)	Biocon Biologics	• CD • PsA (6+) • PsO (6+) • UC	IV infusion: HCP-administered SC Injection: • Self-administered	IV: • 130 mg/26 mg SDV SC: • 45 mg/0.5 mL PFS • 45 mg/0.5 mL SDV • 90 mg/mL PFS	• Refrigerate 36°F to 46°F (2°C to 8°C) • Do not freeze • Do not shake • Store in original carton to protect from light	CD Induction: Vomiting CD: Maintenance: Nasopharyngitis, injection site erythema, vulvovaginal candidiasis/ mycotic infection, bronchitis, pruritus, urinary tract infections, and sinusitis PsO: Nasopharyngitis, upper respiratory tract infection, headache and fatigue UC Induction: Nasopharyngitis UC Maintenance: Nasopharyngitis, headache, abdominal pain, influenza, fever, diarrhea, sinusitis, fatigue and nausea
Yuflyma (adalimumab-aaty)	Celltrion	• AS • CD (6+) • HS • PJIA (2+) • PsO • PsA • RA • UC • UV	SC injection: • Self-administered	SC: • 20 mg/0.2 mL PFS • 40 mg/0.4 mL PFS • 40 mg/0.4 mL PEN • 80 mg/0.8 mL PEN	• Refrigerate at 36°F to 46°F (2°C to 8°C) • Do not freeze • Protect from light	Infections (e.g., upper respiratory, sinusitis), injection site reactions, headache and rash

Autoimmune (AI) Drug List

Select FDA-approved AI specialty medications and infusion therapy

Drug name	Manufacturer	FDA-Approved Indication(s)	Route of Administration	Supply Method	Storage	Select Common Adverse Reactions
Zeposia (ozanimod)	Bristol Myers Squibb	• UC	Oral	Oral: • 7 Day starter pack: – 0.23 mg capsules – 0.46 mg capsules – 0.92 mg capsules • 0.92 mg capsules	• Store at 68°F to 77°F (20°C to 25°C)	Liver test increased, upper respiratory infection and headache



Visit cvs.co/ai-list or scan the QR code to view electronically.
For easy access, you can bookmark this page.

FDA (U.S. Food and Drug Administration). SC (Subcutaneous). IV (Intravenous). PFS (Prefilled syringe). OBI (On-body injector).

AA (Alopecia areata). AD (Atopic dermatitis). aGVHD (Acute graft versus host disease). AOSD (Adult-onset Still's Disease). AS (Ankylosing spondylitis). CAPS (Cryopyrin-associated periodic syndrome). CRS (Cytokine release syndrome). COPD (Chronic obstructive pulmonary disease). CRSwNP (Chronic rhinosinusitis with nasal polyposis). DIRA (Deficiency of IL-1-receptor antagonist). EoE (Eosinophilic esophagitis). ERA (Enthesitis-related arthritis). FCAS (Familial cold autoinflammatory syndrome). FMF (Familial Mediterranean fever). GCA (Giant cell arteritis). HIDS/MKD (Hyperimmunoglobulin D syndrome / mevalonate kinase deficiency). HS (Hidradenitis Suppurativa). JIA (Juvenile idiopathic arthritis). LN (Lupus Nephritis). MWS (Muckle-Wells syndrome). NOMID (Neonatal-onset multisystem inflammatory disease). Nr-axSpA (Non-radiographic axial spondyloarthritis). PJIA (Polyarticular Juvenile idiopathic arthritis). PMR (Polymyalgia rheumatica). PN (Prurigo nodularis). PsA (Psoriatic arthritis). PsO (Plaque psoriasis). RA (Rheumatoid arthritis). RP (Recurrent pericarditis). SJIA (Systemic Juvenile idiopathic arthritis). SLE (Systemic lupus erythematosus). SSc-ILD (Systemic sclerosis - Interstitial lung disease). TRAPS (Tumor necrosis factor receptor-associated periodic syndrome). UC (Ulcerative colitis). UV (Uveitis).

The information contained in this communication is provided in summary form. It is not intended for use as the sole basis of clinical treatment, as a substitute for reading the original research, nor as a substitute for the knowledge, skill and judgment of the medical provider. This list is subject to change. CVS Specialty assumes no liability whatsoever for the information provided or for any diagnosis or treatment made as a result of this information. The information noted in this brochure was taken from the manufacturers' prescribing information. It does not contain all information needed to use these products safely and effectively. See the manufacturers' package insert for full prescribing information. CVS Specialty does not provide medical advice and does not recommend a particular treatment or brand. This document contains references to brand-name prescription drugs that are trademarks or registered trademarks of pharmaceutical manufacturers that are not affiliated with CVS Specialty.